

and wherein said step of extrapolating a withdrawal interval is carried out by:  
determining a half-life multiplier from said first withdrawal time and said residue  
half-life;

determining a concentration at time zero for said first dose from said tolerance  
concentration and said half-life multiplier;

determining a concentration at time zero for said second dose from said first dose,  
said second dose, and said concentration at time zero for said first dose; and then

calculating said withdrawal interval from (a) said residue half-life, (b) said  
concentration at time zero for said second dose, and (c) said tolerance concentration.

*K*  
~~25 30~~  
~~52~~. A method of estimating a withdrawal interval for an adjusted dose of a  
compound from a prior withdrawal time for a corresponding prior dose of said  
compound, corresponding prior half-life data and a tolerance concentration, for a tissue  
of interest, said method comprising the following steps that are performed in a data  
processing system:

accepting selection of an adjusted dose for said compound for which a withdrawal  
interval is to be determined;

extrapolating a withdrawal interval from (a) said prior dose, (b) said prior  
withdrawal time, (c) said half-life data, and (d) said tolerance concentration;

wherein said tolerance concentration is a provisional acceptable residue  
determined by the method comprising:

providing an acceptable daily intake for said compound;

partitioning said acceptable daily intake among tissues according to a set of  
partitioning instructions; and

deriving said provisional acceptable residue for said tissue of interest from  
said partitioned acceptable daily intake.

~~26 31~~  
~~53~~. A method of estimating a withdrawal interval for an adjusted dose of a  
compound from a prior withdrawal time for a corresponding prior dose of said

compound, corresponding prior half-life data and a tolerance concentration, for a tissue of interest, said method comprising the following steps that are performed in a data processing system:

accepting selection of an adjusted dose for said compound for which a withdrawal interval is to be determined;

extrapolating a withdrawal interval from (a) said prior dose, (b) said prior withdrawal time, (c) said half-life data, and (d) said tolerance concentration;

wherein said adjusted dose is modified from said prior dose for species differences, disease differences or both.

19. 32.  
54. A data processing system for estimating a withdrawal interval for an adjusted dose of a compound from a prior withdrawal time for a corresponding prior dose of said compound, corresponding half-life data and a tolerance concentration, for a tissue of interest, said data processing system comprising:

means for accepting selection of an adjusted dose for said compound for which a withdrawal interval is to be determined;

means for extrapolating a withdrawal interval from (a) said prior dose, (b) said prior withdrawal time, (c) said half-life data, and (d) said tolerance concentration;

wherein said half-life data is an empirically determined effective residue half-life, and wherein said means for extrapolating includes:

means for determining a half-life multiplier from said first withdrawal time and said residue half-life;

means for determining a concentration at time zero for said first dose from said tolerance concentration and said half-life multiplier;

means for determining a concentration at time zero for said second dose from said first dose, said second dose, and said concentration at time zero for said first dose; and

means for calculating said withdrawal interval from (a) said residue half-life, (b) said concentration at time zero for said second dose, and (c) said tolerance concentration.